



HCSP FACT SHEET

HCV ADVOCATE

• GENERAL INFORMATION •

Similarities and Differences between HIV and HCV

Written by: Alan Franciscus, Editor-in-Chief

Foreword

HIV and the hepatitis C virus (HCV) share many of the same characteristics, but there are also some very distinct differences in the way they are transmitted, how long each virus lives outside of the body, disease progression, and treatment. This fact sheet will explore some of the similarities and differences between HIV and hepatitis C.

RNA Viruses

Both HIV and hepatitis C are RNA viruses, but they are different types of RNA viruses—HIV is a retrovirus and HCV is a flavivirus.

Replication

HIV mainly infects human immune cells (CD4, macrophages, and dendritic cells). The hepatitis C virus mainly infects liver cells.

The HIV virus goes through a complex process where the HIV virus inserts its genetic material into the host DNA cell and uses the host genetic material to replicate. HCV on the other hand uses the host's cell to replicate, but does not insert itself into the host DNA. These differences have major implications when discussing HIV and HCV treatment and disease progression.

In regards to how much each virus replicates on a daily basis—HIV replicates billions of new viruses vs. the trillions that the hepatitis C virus replicates daily.

Mutation Rate

HIV has a high mutation rate, but hepatitis C is thought to replicate and mutate at a much higher rate than HIV.

Vaccines

Studies are underway to find a vaccine for HIV and hepatitis C, but because of the high mutation rate of both viruses it will most likely be a long time (if ever)

HCSP FACT SHEET

*A publication of the
Hepatitis C Support Project*

EXECUTIVE DIRECTOR, EDITOR-IN-CHIEF, HCSP PUBLICATIONS

Alan Franciscus

DESIGN

*Leslie Hoex,
Blue Kangaroo Design*

PRODUCTION

C.D. Mazoff, PhD

CONTACT INFORMATION

Hepatitis C Support Project
PO Box 427037
San Francisco, CA 94142-7037
alanfranciscus@hcvadvocate.org

The information in this fact sheet is designed to help you understand and manage HCV and is not intended as medical advice. All persons with HCV should consult a medical practitioner for diagnosis and treatment of HCV.

This information is provided by the Hepatitis C Support Project a nonprofit organization for HCV education, support and advocacy

Reprint permission is granted and encouraged with credit to the Hepatitis C Support Project.

Similarities and Differences

before a vaccine that will prevent HIV or hepatitis C infection is discovered and available for use.

Prevalence

In the United States it is estimated that more than 4 million Americans are infected with chronic hepatitis C compared to about 1.2 million who are estimated to be infected with HIV. Worldwide the estimates are in the range of 40 million people infected with HIV compared to 170 million infected with HCV.

Strains

There are different strains of HIV (HIV-1 and HIV-2) and hepatitis C (genotypes 1, 2, 3, 4, 5 and 6). In the United States HIV-1 is the most common strain of HIV and genotype 1 is the most common strain of hepatitis C.

Viral Loads

In people with HIV the viral load affects transmission and disease progression. A high viral load for someone with HIV is in the range of 5,000 to 10,000 copies/mL. In contrast a high HCV viral load for people with hepatitis C is any amount higher than 800,000. Frequently people with HCV have viral load amounts in the millions. Studies have not found that a high hepatitis C viral load increases transmission of hepatitis C or HCV disease progression. However, there is some evidence that a high HCV viral load increases the chances of transmission of HCV from a mother to child.

How long do HIV and HCV live outside the body?

Once HIV is exposed to air it lasts only a few moments, but may live longer if blood is present especially in a closed environment such as in a syringe. Hepatitis C on the other hand has been found to live outside the body for at least 16 hours but no longer than 4 days, and in a syringe for up to 63 days. This is the reason why, for instance, that exposure to blood

through unsafe tattooing carries a risk of transmission of hepatitis C, but it does not carry a risk of transmission of HIV.

Chronic Infection

All people who are initially infected with HIV will have long-term or chronic infection. Hepatitis C is different—some people (about 20 to 40%) who are initially infected (acute) will naturally resolve or clear HCV infection. The other 60 to 80% of people will have long-term or chronic infection.

Transmission

HIV and hepatitis C are transmitted in similar ways through blood routes although there are clear differences in the risk of transmission. The hepatitis C virus is about 10 times more concentrated in blood than HIV. This means that the hepatitis C virus can be transmitted easier when any HCV-infected blood is involved.

In the United States, the most common way that HIV is transmitted is through anal and vaginal sex. HIV can also be transmitted by sharing HIV infected needles and an HIV positive mother can pass HIV to her child.

The most common way that hepatitis C is transmitted is from sharing HCV-infected needles and drug preparation and using tools. Transmission routes of hepatitis C that are less common but still possible include sexual, and mother-to-child.

Treatment

Hepatitis C can be cured—this is because the hepatitis C virus does not integrate into the host's DNA when it is replicating (see above). It's a different story with HIV because HIV inserts itself into the host's DNA. As a result, HIV treatment is aimed at lowering HIV viral load, increasing or stabilizing CD4 cell counts and preventing long-term consequences of infection.

Similarities and Differences

Drug Resistance

The medications used to treat HIV can lead to drug resistance because they are direct antiviral medications. The newly approved protease inhibitors used in combination with pegylated interferon and ribavirin for people infected with chronic HCV genotype 1 can also lead to drug resistance.

People infected with HCV genotypes 2 or 3 are only prescribed pegylated interferon and ribavirin—these two medications do not lead to drug resistance.

Thank you to Heather Lusk for review and input.

Related publications:

- **Easy C - A Guide to HIV and Hep C Coinfection**
www.hcvadvocate.org/hepatitis/easyfacts/Easy%20Coinfec%20GD.pdf
- **Frequently Asked Questions about HIV & Hepatitis C**
www.hcvadvocate.org/hepatitis/factsheets_pdf/Coinfection_FAQ.pdf
- **A Guide to HIV/HCV and Hep C Coinfection**
www.hcvadvocate.org/hepatitis/factsheets_pdf/Coinfection%20Guide.pdf

For more information

- | | |
|--|--|
| <ul style="list-style-type: none">• Centers for Disease Control and Prevention
www.cdc.gov• HIV and Hepatitis.Com
www.hivandhepatitis.com | <ul style="list-style-type: none">• National AIDS Treatment Advocacy Project:
www.natap.org• Project Inform
www.projectinform.org |
|--|--|

Visit our websites to learn more about viral hepatitis:

www.hcvadvocate.org • www.hbvadvocate.org
www.hepatitistattoos.org

